



NOTICE: AVAILABILITY OF LOPINAVIR/RITONAVIR ORAL PELLETS TO CHILDREN ON ANTIRETROVIRAL TREATMENT

The Paediatric Hospital Level Standard Treatment Guidelines (STGs) and Essential Medicines List (EML) currently recommends the use of lopinavir/ritonavir (LPV/r) as part of the first-line antiretroviral treatment regimen in children from 1 month of age¹. This is currently supplied as a solution (80/20mg/ml). The poor palatability of the solution may result in spitting out, vomiting or refusal of medication and thereby affect adherence to treatment. LPV/r pellets (40/10mg per capsule) is recommended for children over the age of 6 months, who cannot tolerate the LPV/r solution. This formulation requires initiation by a doctor. See Table 1 below for dose recommendations according to weight:

Table 1: Dosing of LPV/r pellets according to weight

| Lopinavir/ritonavir (LPV/r) Oral Pellets <i>Only for use in patients not tolerating LPV/r solution</i> | |
|--|--|
| Target dose | 300/75 mg/m ² /dose LPV/r TWICE daily |
| Note | Pellets 40/10 mg per capsule CAPSULES ARE NOT RECOMMENDED <6 MONTHS OF AGE CAPSULES MUST NOT BE SWALLOWED WHOLE |
| <i>Weight</i> | |
| 3-4.9 | 2 capsules twice daily |
| 5-5.9 | 2 capsules twice daily |
| 6-9.9 | 3 capsules twice daily |
| 10-13.9 | 4 capsules twice daily |
| 14-19.9 | 5 capsules twice daily |
| 20-24.9 | 6 capsules twice daily |
| 25-29.9 | 7 capsules twice daily |
| 30-34.9 | 8 capsules twice daily |
| ≥35 | 10 capsules twice daily |

The instructions on how to administer LPV/r pellets to children are as follows:

- **Do not swallow capsule whole.**
- Hold the capsule at both ends and, twisting in opposite directions, and then pull apart to pour out the pellets.
- Add the pellets (from the required number of tablets) to a spoonful of food a little at a time. For example, porridge can be used (not too hot, must be at room temperature).
- Do not stir, crush or dissolve the pellets; rather sprinkle over the food.
- Use only a small amount of food, to ensure child can consume all the pellets (never keep food with pellets in for later use). Discard food with pellets after 2 hours.
- The capsule cover can be discarded with usual waste.

¹ Standard Treatment Guidelines and Essential Medicines List for South Africa. Hospital Level: Paediatrics, 2017 Edition

NOTE: Once the child can swallow tablets, they should be placed on the LPV/r 100/25mg or 200/50mg tablet according to the weight-based dosing table². For recommended regimens, please refer to the 2019 ART Clinical Guidelines for the Management of HIV in Adults, Pregnancy, Adolescents, Children, Infants and Neonates, October 2019².

For further assistance on how and when to use this formulation please contact:

National HIV & TB Care Worker Hotline: 0800 212 506 or 021 406 6782

National HIV & TB Care Worker Hotline: 071 840 1572

- This helpline can be contacted *via* SMS/Please call me/WhatsApp

Right to Care Paediatric and Adolescent HIV Helpline 082 352 6642

Right to Care Adult HIV Helpline 082 957 6698

- These helplines can be contacted via call/SMS/Please call me/WhatsApp

KZN Paediatric Hotline 0800 006 603

Procurement of LPV/r pellets

Public healthcare facilities may procure LPV/r pellets (40/10mg per capsule) on quotation as a buy-out item (see Annexure A: Lopimune ® 40/10mg pellets - quotation).

Provinces and Healthcare facilities are requested to distribute and communicate this information in consultation with their Pharmaceutical and Therapeutics Committees. Kindly share with all healthcare professionals and relevant stakeholders.

Comments may be submitted *via* e-mail:

Stock queries:

Ms Maureen Masinge

Tel: 012 395 9683

E-mail: Maureen.Masinge@health.gov.za


Clinical queries:

Essential Drugs Programme

E-mail: SAEDP@health.gov.za

The National Department of Health continues to explore other therapeutic options and formulations to improve access to antiretroviral treatment.

Kind regards



MS K JAMALOODIEN

DIRECTOR: AFFORDABLE MEDICINES

DATE: 1/4/2020

² Available online at <https://www.knowledgehub.org.za/elibrary/2019-art-clinical-guidelines-management-hiv-adults-pregnancy-adolescents-children-infants>